

JOURNAL OF NUTRITIONAL SCIENCE AND VITAMINOLOGY

Vol 64, No 4, August 2018

CONTENTS

Regular Paper

- IKEDA S, NOMURA S, HANZAWA F, TAKAHASHI S, ODA H, FUJIWARA Y and UCHIDA T: α -Tocopherol Intake Decreases Phylloquinone Concentration in Bone but Does Not Affect Bone Metabolism in Rats 243–250
- TAKASAKI M, MOMOSAKI R, WAKABAYASHI H and NISHIOKA S: Construction and Quality Evaluation of the Japanese Rehabilitation Nutrition Database 251–257
- IIZUKA Y, OZEKI A, TANI T and TSUDA T: Blackcurrant Extract Ameliorates Hyperglycemia in Type 2 Diabetic Mice in Association with Increased Basal Secretion of Glucagon-Like Peptide-1 and Activation of AMP-Activated Protein Kinase 258–264
- FUKUNAGA S, WADA S, SATO T, HAMAGUCHI M, AOI W and HIGASHI A: Effect of *Torula* Yeast (*Candida utilis*)-Derived Glucosylceramide on Skin Dryness and Other Skin Conditions in Winter 265–270
- HASHIMOTO K, ZAIMA N, SEKIGUCHI H, KUGO H, MIYAMOTO C, HOSHINO K, KAWASAKI N, SUTOH K, USUMI K and MORIYAMA T: Dietary DNA Attenuates the Degradation of Elastin Fibers in the Aortic Wall in Nicotine-Administrated Mice 271–276
- NISHIJIMA C, SATO Y, CHIBA T and UMEGAKI K: Nationwide Online Survey to Complement the Current Voluntary Reporting System for Adverse Events Associated with Dietary Supplements: Application to the Case of Skin Manifestations 277–283

Note

- TANAKA H, NISHIKAWA Y, KURE K, TSUDA K and HOSOKAWA M: The Addition of Xanthan Gum to Enteral Nutrition Suppresses Postprandial Glycemia in Humans 284–286
- TSUCHIYA Y and KAWAMATA K: [6]-Gingerol Induces Amiloride-Sensitive Sodium Absorption in the Rat Colon via the Capsaicin Receptor TRPV1 in Colonic Mucosa 287–291
- SHIBATA K: Urinary Excretion of 2-Oxo Acids Is Greater in Rats with Streptozotocin-Induced Diabetes 292–295
- SANEYASU T, SHINDO H, HONDA K and KAMISOYAMA H: The Extract of Soybean Protein Increases Slow-Myosin Heavy Chain Expression in C2C12 Myotubes 296–300
- ASHIGAI H, KOMANO Y, WANG G, KAWACHI Y, SUNAGA K, YAMAMOTO R, TAKATA R and YANAI T: Orally Administered Polysaccharide Derived from Blackcurrants (*Ribes nigrum* L.) Improves Skin Hydration in Ultraviolet-Irradiated Hairless Mice 301–304

Withdrawal 305